


Europ. J. Combinatorics (2001) **22**, 513–519

doi:10.1006/eujc.2000.0413

Available online at <http://www.idealibrary.com> on  IDEAL[®]

Suzuki Groups and 2- $(v, k, 1)$ Designs

WEIJUN LIU, HUILING LI AND CHUNGUI MA

If one lets \mathcal{D} be a 2- $(v, k, 1)$ design with $G \leq \text{Aut}(\mathcal{D})$ block-primitive, then G does not have a Suzuki group as its socle.

© 2001 Academic Press

1. INTRODUCTION

A 2- $(v, k, 1)$ design $\mathcal{D} = (\Omega, \mathcal{B})$ is a system consisting of a finite set Ω of v points and a collection \mathcal{B} of k -subsets of Ω , called blocks, such that any 2-subset of Ω is contained in precisely one block. We assume throughout that $2 < k < v$.

Let $G \leq \text{Aut}(\mathcal{D})$ be a group of automorphisms of a 2- $(v, k, 1)$ design \mathcal{D} . The group G is said to be *block-transitive* (*block-primitive*, respectively) on \mathcal{D} if G is transitive (primitive, respectively) on \mathcal{B} . Group G is said to be *point-transitive* (*point-primitive*, respectively) on \mathcal{D} if G is transitive (primitive, respectively) on Ω . A *flag* of \mathcal{D} is a pair consisting of a point and a block containing this point. G is *flag-transitive* on \mathcal{D} if G is transitive on the set of flags of \mathcal{D} . The following results are well known: if G is block transitive, then G is also point-transitive (see [1]); if G is flag-transitive, then G is point-primitive (see [8]); if G acts block primitively on a finite projective plane, then G is point-primitive (see [11]). Doyen and Delandtsheer [5] conjecture that if G is block-primitive, then G is also point-primitive.

It is known that the conjecture is true under any of the following hypotheses:

- (1) \mathcal{D} is a finite projective plane;
- (2) $k/(k, v) \leq 4$;
- (3) $v > \frac{[k(k-1)/2-1]^2}{2}$;
- (4) $k \leq 40$;
- (5) G has a subgroup acting regularly on Ω ;
- (6) the rank of G acting on \mathcal{B} does not exceed 7.

Results (1), (3), (5) and (6) can be found in [5]; (2) and (4) in [5, 7, 15]. Delandtsheer [6] proved the following result: If one lets \mathcal{D} be a 2- $(v, k, 1)$ design other than a projective plane and $G \leq \text{Aut}(\mathcal{D})$ be block-primitive, then G is almost simple, i.e., there is a non-abelian simple T such that $T \trianglelefteq G \leq \text{Aut}(T)$. Li and Liu proved the conjecture holds if $T = \text{soc}(G)$ is an alternating group A_n or a sporadic simple group, respectively. In [17], Zhou *et al.* proved the above conjecture is true if $T \cong {}^2G_2(q)$.

In this paper, we prove the following theorem.

THEOREM 1.1. *If one lets \mathcal{D} be a 2- $(v, k, 1)$ design and $G \leq \text{Aut}(\mathcal{D})$ be block-primitive, then the socle of G is not isomorphic to $Sz(q)$.*

Thus this supports Doyen–Delandtsheer’s conjecture.

The second section describes the notation and contains a lot of preliminary results regarding the simple Suzuki groups $Sz(q)$ and 2- $(v, k, 1)$ designs.

In the third section, we will give the proof of Theorem 1.1.

2. PRELIMINARY RESULTS

In this section, we begin by restating some fundamental properties of the simple Suzuki groups $Sz(q)$.

We put $q = 2^{2m+1}$, $t = 2^m$ and $G = Sz(q)$. By Theorem 3.10 of Chapter XI of [9], G possesses cyclic subgroups U_1 and U_2 of orders $q + 2t + 1$ and $q - 2t + 1$, respectively. U_1 and U_2 are Hall subgroups of G . By Lemma 3.1 of Chapter XI of [9], G possesses subgroups F and H , where F is a 2-group of exponent 4, class 2 and order q^2 . H is isomorphic to $GF(q)^\times$, a multiplicative group of $GF(q)$.

LEMMA 2.1 ([9, CHAPTER XI, LEMMA 3.12]). *Every maximal subgroup of G is conjugate to one of the following:*

- (1) $Sz(a)$, $a^i = 2^{2m+1}$, i a prime;
- (2) FH ;
- (3) $N_G(H)$;
- (4) $N_G(U_i)$, $i = 1, 2$.

Conversely, there is exactly one class of maximal subgroups of G for each case in Lemma 2.1. Since $|Sz(q)| = q^2(q^2 + 1)(q - 1)$ and $q^2 + 1 = (q + 2t + 1)(q - 2t + 1)$, and q^2 , $q + 2t + 1$, $q - 2t + 1$, $q - 1$ are mutually prime, the Sylow p -subgroups of $Sz(q)$ are conjugate to the subgroups of F , H , U_1 or U_2 .

LEMMA 2.2 ([9, CHAPTER XI]). *For any $g \in G$, we have:*

- (1) if $F \neq F^g$, then $F \cap F^g = \{1\}$;
- (2) if $H \neq H^g$, then $H \cap H^g = \{1\}$, and for all $1 \neq h \in H$, $C_G(h) = H$;
- (3) if $U_i \neq U_i^g$, then $U_i \cap U_i^g = \{1\}$, and for all $1 \neq u \in U_i$, $C_G(u) = U_i$, where $i = 1$ and 2.

LEMMA 2.3. *Let P be a Sylow p -subgroup of G :*

- (1) if $P = F$, then $N_G(P) = FH$;
- (2) if $P \leq H$, then $N_G(P) = N_G(H)$;
- (3) if $P \leq U_i$, then $N_G(P) = N_G(U_i)$, where $i = 1$ and 2.

PROOF. (1) By Chapter XI, Lemma 3.1 of [9], $H \leq N_G(F)$, thus $FH \leq N_G(F)$. Since FH is a maximal subgroup of G , it follows that $N_G(F) = FH$.

(2) For any $g \in N_G(P)$, $P \leq H \cap H^g$. By Lemma 2.2, $H = H^g$, this means $g \in N_G(H)$. Thus $N_G(P) \leq N_G(H)$. Since the group H is cyclic, any element of G which normalizes H will normalize any Sylow p -subgroup of H . Thus $N_G(P) = N_G(H)$.

(3) The proof is similar to that of (2).

Let \mathcal{D} be a 2- $(v, k, 1)$ design. Ω is the set of points of \mathcal{D} , \mathcal{B} is the set of blocks of \mathcal{D} , b is the number of blocks of \mathcal{D} and r is the number of blocks of \mathcal{D} through a given point of Ω . Then

$$b = \frac{v(v-1)}{k(k-1)}, \quad r = \frac{(v-1)}{(k-1)}.$$

Let

$$b_1 = (b, v), \quad b_2 = (b, v-1), \quad k_1 = (k, v) \quad \text{and} \quad k_2 = (k, v-1).$$

Obviously,

$$k = k_1 k_2, \quad b = b_1 b_2, \quad r = b_2 k_2 \quad \text{and} \quad v = b_1 k_1.$$

Let B be a block of \mathcal{D} . Then, G_B will be the block stabilizer and $G_{(B)}$ the pointwise stabilizer of the block.

If G is a block-transitive automorphism group of \mathcal{D} , and not primitive on Ω . We define \mathcal{C} as follow: \mathcal{C} is a set of some non-trivial partition of Ω which is preserved by G and on which G acts primitively. Let c be the cardinality of \mathcal{C} and d the common size of the classes in \mathcal{C} . Therefore, $v = cd$. \square

LEMMA 2.4. (1) (Lemma 2 of [7]). There exist positive integers x and y , such that $c = xb_2 + 1$ and $d = yb_2 + 1$.

(2) (Lemma 4 of [6]). There exists a prime number p dividing b but not v , and $b > v$.

The following Lemmas are very useful for our proof of Theorem 1.1.

LEMMA 2.5 (LEMMA 2 OF [4]). Let G act as a block-transitive automorphism group of a $2-(v, k, 1)$ design. Let B be a block and H a subgroup of G_B . Assume that H satisfies the two following conditions:

- (i) $|\text{Fix}(H) \cap B| \geq 2$ and;
- (ii) if $K \leq G_B$ and $|\text{Fix}(K) \cap B| \geq 2$ and K is conjugate to H in G then H is conjugate to K in G_B .

Then either (a) $\text{Fix}(H) \subseteq B$ or (b) the induced structure on $\text{Fix}(H)$ is a $2-(v_0, k_0, 1)$ design where $v_0 = |\text{Fix}(H)|$, $k_0 = |\text{Fix}(H) \cap B|$. Furthermore, $N_G(H)$ acts as a block-transitive group on this design.

LEMMA 2.6. Let G act as a block-transitive automorphism group of a $2-(v, k, 1)$ design. Let B be a block and v even. Assume that there exists a 2-subgroup P of G_B such that $\text{Fix}(P) \subseteq B$. Then $k \mid v$ and G is flag-transitive.

PROOF. Let i be an involution in P . As v is even, then k is even. For any block $B' \neq B$, if $B^i = B'$, then B' has no point fixed by i . Suppose B_1 and B_2 are two blocks fixed by i . Then $B_1 \cap B_2 = \emptyset$. In fact, if $\alpha \in B_1 \cap B_2$, then α is fixed by i . Hence i fixes at least one other point of B_1 , and this implies that $B = B_1$. Similarly, we may derive $B = B_2$. Consider the cycle decomposition of i acting on Ω , we have $|\text{Fix}_B(\langle i \rangle)| = (v - k)/k + 1 = v/k$, this means $k \mid v$, where $\text{Fix}_B(\langle i \rangle)$ denotes the set of blocks fixed by i . By [3], G is flag-transitive. \square

LEMMA 2.7. Let G act as a block-transitive automorphism group of a $2-(v, k, 1)$ design. Let B be a block and let i be an involution of G_B . Assume that G_B has a unique conjugate class of involutions. If $|\text{Fix}(\langle i \rangle) \cap B| \geq 2$ and v is even, then G is flag-transitive or the induced structure on $\text{Fix}(\langle i \rangle)$ is a $2-(v_0, k_0, 1)$ design where $v_0 = |\text{Fix}(\langle i \rangle)|$, $k_0 = |\text{Fix}(\langle i \rangle) \cap B|$. Furthermore, $N_G(\langle i \rangle)$ acts as a block-transitive group on this design.

PROOF. Set $H = \langle i \rangle$. Clearly H satisfies the hypothesis of Lemma 2.5. If $\text{Fix}(H) \subseteq B$ then G is flag transitive by Lemma 2.6. If $\text{Fix}(H) \not\subseteq B$, then by Lemma 2.5, Lemma 2.7 is true. \square

LEMMA 2.8. Let $G \leq \text{Aut}(\mathcal{D})$, G act block-transitively on \mathcal{D} . If there exists a prime number p such that $p \mid b$ but $p \nmid v$, then for some $\alpha \in \Omega$, $N_G(P) \leq G_\alpha$, where P is a Sylow p -subgroup of G .

PROOF. (i) P does not fix any block of \mathcal{D} .

Since $b = |G|/|G_B|$ and $p|b$, it follows that $P \not\leq G_B$ for any $B \in \mathcal{B}$.

(ii) P fixes exactly one point of Ω .

Suppose that P does not fix any point of Ω , then for any point β of Ω , $|\beta^P| = |P : P_\beta| > 1$. Since $v = \sum |\beta^P|$, $p|v$. This conflicts with the hypothesis. Hence P fixes exactly one point by (i).

(iii) $N_G(P) \leq G_\alpha$.

For any $g \in N_G(P)$, $\alpha^{g^P} = \alpha^{P^g} = \alpha^g$. So α^g is fixed by P . By (ii), $\alpha = \alpha^g$, thus $N_G(P) \leq G_\alpha$. \square

LEMMA 2.9 ([12, 13]). *Let G be a finite group and M a maximal subgroup of G not containing $\text{Soc}(G)$. If $\text{Soc}(G) = \text{Sz}(q)$, where $q = 2^{2m+1}$ and $m > 0$, then $M \cap \text{Sz}(q)$ is a maximal subgroup of $\text{Sz}(q)$.*

In fact, it is not difficult to prove the above lemma.

3. PROOF OF THEOREM 1.1

First we prove the following proposition.

PROPOSITION 3.1. *Let \mathcal{D} be a 2 -($v, k, 1$) design and $G \leq \text{Aut}(\mathcal{D})$ be block-primitive. If $\text{Sz}(q) \trianglelefteq G \leq \text{Aut}(\text{Sz}(q))$, then G is also point-primitive.*

PROOF. Since G is block-primitive, we have that G_B is a maximal subgroup of G for any block B of \mathcal{D} . By $\text{Sz}(q) \trianglelefteq G$ we get $\text{Sz}(q)$ is block-transitive. Thus $G_B \cap \text{Sz}(q) \neq \text{Sz}(q)$. By Lemma 2.9, we may assume that $G = \text{Sz}(q)$.

Suppose that there exist counterexamples (\mathcal{D}, G) . Then there is a non-trivial partition $\mathcal{C} = \{C = C_1, C_2, \dots, C_c\}$, $|C_i| = d$, $i = 1, 2, \dots, c$, such that G acts primitively on \mathcal{C} , and this action is faithful (see [6]). Thus G_C is a maximal subgroup of G and G_C is one of the groups of Lemma 2.1. By Lemma 2.4, there is a prime p , such that $p|b$ but $p \nmid v$. Hence by Lemma 2.8 there exists a Sylow p -subgroup P of G , such that $N_G(P) \leq G_\alpha < G_C$.

(1) If $G_C \cong \text{Sz}(a)$, $q = a^i$ and i is a prime, then by Lemmas 2.3 and 2.8, G_C contains H or U_1 or U_2 . This is impossible.

(2) If $G_C \cong FH$, then $p = 2$ or p divides $q - 1$. Thus $N_G(P) \cong FH$ or $N_G(H)$ by Lemma 2.3. Since FH and $N_G(H)$ are the maximal subgroups of G , we have $G_\alpha = G_C$, a contradiction.

(3) If $G_C \cong N_G(H)$, then p divides $q - 1$. By Lemma 2.3, we have $N_G(P) = N_G(H)$ and hence $G_\alpha = G_C$. Again we get a contradiction.

(4) If $G_C \cong N_G(U_i)$, where $i = 1, 2$, then p divides $q + (-1)^i 2t + 1$ and $P \leq U_i$. Therefore $N_G(P) = N_G(U_i)$ by Lemma 2.3, which conflicts with $G_\alpha < G_C$.

By (1)–(4) above we conclude that G_α is a maximal subgroup of G . Hence G acts primitively on Ω and so does G , which gives rise to a contradiction. Thus the assertion of the proposition holds. \square

Now we start to prove our main theorem.

Suppose $\text{Soc}(G) \cong \text{Sz}(q)$, then by Lemma 2.9, we may assume that $G = \text{Sz}(q)$. Thus G_α and G_B are all maximal subgroups of G and so they occur in Lemma 2.1. Clearly \mathcal{D} is not a projective plane (see [11]). It follows that $b > v$, i.e., $|G_B| < |G_\alpha|$. By [10], $G_\alpha \not\cong FH$. If $G_\alpha \cong \text{Sz}(a)$, where a satisfies $q = a^m$, m a prime, then by Lemma 2.8, there always exists a

Sylow p -subgroup P of G , where p is odd, such that $N_G(P) \leq G_\alpha$. According to Lemma 2.3, G_α contains subgroups H or U_1 or U_2 of G . This is impossible. Thus $G_\alpha \not\cong Sz(a)$.

First we eliminate the case $G_B \cong Sz(a)$.

In this case, $G_\alpha \cong N_G(U_1)$, $N_G(U_2)$ or $N_G(H)$. In order to calculate $|\text{Fix}(\langle i \rangle)|$, we consider the pair (i, G_α) , where i is an involution of G such that i is contained in G_α . Since G has a unique conjugacy class of involutions (see [16]) by counting pairs on two ways we see that

$$|G : G_\alpha| \cdot e(G_\alpha) = e(G) \cdot N,$$

where $e(G)$ denotes the number of involutions of G , and N denotes the number of conjugates of G_α which contains i , it is equal to the number of points fixed by i , i.e., $N = |\text{Fix}(\langle i \rangle)|$. Hence

$$|\text{Fix}(\langle i \rangle)| = \frac{|G : G_\alpha| \cdot e(G_\alpha)}{e(G)}.$$

By [16],

$$e(G) = |G : C_G(i)| = (q^2 + 1) \cdot (q - 1).$$

Thus

$$|\text{Fix}(\langle i \rangle)| = \frac{q^2 \cdot e(G_\alpha)}{|G_\alpha|}. \quad (*)$$

By [16],

$$N_G(U_1) = \langle u, s \mid o(u) = q + 2t + 1, \quad o(s) = 4, \quad u^s = u^q \rangle.$$

$s^2 u^j$ are involutions of $N_G(U_1)$, where $j = 1, 2, \dots, q + 2t + 1$. Thus $e(N_G(U_1)) = q + 2t + 1$. Similarly, $e(N_G(U_2)) = q - 2t + 1$ and $e(N_G(H)) = q - 1$.

If $G_\alpha \cong N_G(U_1)$, by (*), $|\text{Fix}(\langle i \rangle)| = q^2/4$. Set $Q = \langle i \rangle$. By the block transitivity of G , we may assume that $Q \leq G_B$. Clearly G_B and $Q = \langle i \rangle$ satisfy the conditions of Lemma 2.7. Then G is flag transitive or the induced structure on $\text{Fix}(Q)$ is a 2- $(q^2/4, k_0, 1)$ design, where $k_0 = |\text{Fix}(Q) \cap B|$. Furthermore, $N_G(Q)$ acts as a block-transitive on this design. By [2] and [14], G is not flag-transitive. Thus the latter holds. Since $N_G(Q) \cong F$, $|N_G(Q)| = q^2$ (see [16]). It follows that $b_0 \mid q^2$, where b_0 is the number of blocks of the above-induced structure. Since $b_0 \geq v_0 = q^2/4$, $b_0 = q^2, q^2/2$ or $q^2/4$. Thus we have the equality

$$4k_0^2 - 4k_0 = \frac{q^2}{4} - 1$$

or

$$2k_0^2 - 2k_0 = \frac{q^2}{4} - 1$$

or

$$k_0^2 - k_0 = q^2/4 - 1.$$

When $m > 1$, these equalities are not satisfied.

Therefore, when $G_B \cong Sz(a)$, then $G_\alpha \not\cong N_G(U_1)$.

Similarly, we may prove that if $G_B \cong Sz(a)$, then $G_\alpha \not\cong N_G(U_2)$ and $G_\alpha \not\cong N_G(H)$. Thus $G_B \not\cong Sz(a)$.

Next we discuss the remaining cases of G_α and G_B .

(i) $G_\alpha \cong N_G(U_1)$

We have

$$v = \frac{q^2(q^2 + 1)(q - 1)}{4(q + 2t + 1)} = \frac{q^2(q - 2t + 1)(q - 1)}{4}.$$

By $|G_B| < |G_\alpha|$, $G_B \cong N_G(U_2)$ or $N_G(H)$.

If $G_B \cong N_G(U_2)$, then $b_2 = q + 2t + 1$. Since $b_2 k_2(k - 1) = v - 1$, we have

$$(q + 2t + 1)k(k - 1) = (q - 2t + 1) \cdot (q^2(q - 2t + 1)(q - 1)/4 - 1).$$

When $m > 1$, the left-hand side of this equality is even, but the right-hand side is odd, a contradiction.

If $G_B \cong N_G(H)$, then

$$b_2 = \frac{b}{b_1} = \frac{b}{(b, v)} = 2(q + 2t + 1).$$

But then $(b_2, v) \neq 1$. This is a contradiction.

(ii) $G_\alpha \cong N_G(U_2)$

Clearly,

$$v = \frac{q^2(q + 2t + 1)(q - 1)}{4},$$

and $G_B \cong N_G(H)$. Thus $b_2 = 2(q - 2t + 1)$, conflicting with $(b_2, v) = 1$.

(iii) $G_\alpha \cong N_G(H)$

By $|G_B| < |G_\alpha|$, then $G_B \cong Sz(a)$. This conflicts with the statement as above.

To sum up, we have derived a contradiction. Thus G is not isomorphic to $Sz(q)$. This completes the proof of Theorem 1.1.

ACKNOWLEDGEMENTS

Supported by the National Natural Science Foundation of China and the Natural Science Foundation of Zhejiang Province of China. The authors would like to thank the referee for pointing out errors in the original version of this paper.

REFERENCES

1. R. E. Block, On the orbits of collineation groups, *Math. Z.*, **96** (1961), 33–49.
2. F. Buekenhout, A. Delandtsheer, J. Doyen, P. B. Kleidman, M. W. Liebeck and J. Saxl, Linear spaces with flag-transitive automorphism groups, *Geom. Ded.*, **36** (1990), 89–94.
3. A. Camina and T. M. Gagen, Block transitive automorphism groups of designs, *J. Algebra*, **86** (1984), 549–554.
4. A. Camina and J. Siemons, Block transitive automorphism groups of 2 -($v, k, 1$) block designs, *J. Comb. Theory, Ser. A*, **51** (1989), 268–276.
5. A. Delandtsheer, Line-primitive groups of small rank, *Discrete Math.*, **68** (1988), 103–106.
6. A. Delandtsheer, Line-primitive automorphism groups of finite linear spaces, *Europ. J. Combinatorics*, **10** (1989), 161–169.
7. W. Fang and H. Li, A generalization of Camina-Gagen's Theorem, *J. Math., (China)*, **13** (1993), 437–441.
8. D. G. Higman and J. E. McLaughlin, Geometric ABA-groups, *Illinois J. Math.*, **5** (1961), 382–397.
9. B. Huppert and N. Blackburn, *Finite Groups III*, Springer-Verlag, New York, 1982.
10. W. M. Kantor, Homogeneous designs and geometric lattices, *J. Comb. Theory, Ser. A*, **38** (1985), 66–74.
11. W. M. Kantor, Primitive permutation groups of odd degree and application to finite projective plane, *J. Algebra*, **106** (1987), 15–45.
12. P. B. Kleidman, A note on the Suzuki groups $Sz(q)$, unpublished.

13. P. B. Kleidman and M. W. Liebeck, A survey of the maximal subgroups of the finite simple groups, *Geom. Ded.*, **25** (1988), 375–389.
14. M. W. Liebeck, The classification of finite linear spaces with flag-transitive automorphism groups of affine type, *J. Comb. Theory, Ser. A*, **84** (1998), 196–235.
15. W. Liu, Line-primitive 2 -($v, k, 1$) designs with k small, *J. Math. (PRC)*, **15** (1995), 375–380.
16. M. Suzuki, On a class of doubly transitive groups, *Ann. Math.*, **75** (1962), 105–145.
17. S. Zhou, H. Li and W. Liu, The Ree groups ${}^2G_2(q)$ and 2 -($v, k, 1$) block designs, *Discrete Math.*, **224** (2000), 251–258.

Received 11 June 1999 and accepted 30 July 2000

WEIJUN LIU

*Department of Mathematics, Physics and Mechanics,
Changsha Railway University,
Changsha,*

Hunan, 410075,

People's Republic of China

and

*Department of Mathematics,
Zhejiang University,*

Hangzhou,

Zhejiang, 310027,

People's Republic of China

AND

HUILING LI AND CHUNGUI MA

*Department of Mathematics,
Zhejiang University,*

Hangzhou,

Zhejiang, 310027,

People's Republic of China